



# The VVV Survey RR Lyrae Population in the Galactic Center Region\*

Rodrigo Contreras Ramos<sup>1,2</sup> , Dante Minniti<sup>1,3,4</sup> , Felipe Gran<sup>1,2</sup> , Manuela Zoccali<sup>1,2</sup> , Javier Alonso-García<sup>1,5</sup> , Pablo Huijse<sup>1,6</sup> , María Gabriela Navarro<sup>1,3</sup> , Álvaro Rojas-Arriagada<sup>1,2</sup> , and Elena Valenti<sup>7</sup>

<sup>1</sup> Instituto Milenio de Astrofísica, Santiago, Chile; [rcontrer@astro.puc.cl](mailto:rcontrer@astro.puc.cl)

<sup>2</sup> Pontificia Universidad Católica de Chile, Instituto de Astrofísica, Av. Vicuña Mackenna 4860, Santiago, Chile

<sup>3</sup> Departamento de Ciencias Físicas, Facultad de Ciencias Exactas, Universidad Andrés Bello, Av. Fernández Concha 700, Las Condes, Santiago, Chile

<sup>4</sup> Vatican Observatory, V00120 Vatican City State, Italy

<sup>5</sup> Unidad de Astronomía, Facultad Cs. Básicas, Universidad de Antofagasta, Avda. U. de Antofagasta 02800, Antofagasta, Chile

<sup>6</sup> Instituto de Informática, Universidad Austral de Chile, General Lagos 2086, Valdivia, Chile

<sup>7</sup> European Southern Observatory, Karl-Schwarzschild-Str. 2, D-85748 Garching bei Muenchen, Germany

Received 2018 April 6; revised 2018 May 23; accepted 2018 May 25; published 2018 August 13

## Abstract

Deep near-IR images from the VISTA Variables in the Vía Láctea (VVV) Survey were used to search for RR Lyrae stars within 100 arcmin from the Galactic Center. A large sample of 960 RR Lyrae of type ab (RRab) stars were discovered. A catalog is presented featuring the positions, magnitudes, colors, periods, and amplitudes for the sample, in addition to estimated reddenings, distances, and metallicities, and measured individual relative proper motions. We use the reddening-corrected Wesenheit magnitudes, defined as  $W_{K_s} = K_s - 0.428 \times (J - K_s)$ , in order to isolate bona fide RRL belonging to the Galaxy Center, finding that 30 RRab are foreground/background objects. We measure a range of extinctions from  $A_{K_s} = 0.19$  to 1.75 mag for the RRab in this region, finding that large extinction is the main cause of the sample incompleteness. The mean period is  $P = 0.5446 \pm 0.0025$  days, yielding a mean metallicity of  $[\text{Fe}/\text{H}] = -1.30 \pm 0.01$  ( $\sigma = 0.33$ ) dex for the RRab sample in the Galactic Center region. The median distance for the sample is  $D = 8.05 \pm 0.02$  kpc. We measure the RRab surface density using the less reddened region sampled here, finding a density of 1000 RRab/sq deg at a projected Galactocentric distance  $R_G = 1.6$  deg. Under simple assumptions, this implies a large total mass ( $M > 10^9 M_\odot$ ) for the old and metal-poor population contained inside  $R_G$ . We also measure accurate relative proper motions, from which we derive tangential velocity dispersions of  $\sigma V_l = 125.0$  and  $\sigma V_b = 124.1$  km s<sup>-1</sup> along the Galactic longitude and latitude coordinates, respectively. The fact that these quantities are similar indicate that the bulk rotation of the RRab population is negligible, and implies that this population is supported by velocity dispersion. In summary, there are two main conclusions of this study. First, the population as a whole is no different from the outer bulge RRab, predominantly a metal-poor component that is shifted with respect to the Oosterhoff type I population defined by the globular clusters in the halo. Second, the RRab sample, as representative of the old and metal-poor stellar population in the region, has high velocity dispersions and zero rotation, suggesting a formation via dissipational collapse.

**Key words:** catalogs – Galaxy: bulge – proper motions – stars: variables: RR Lyrae – surveys

**Supporting material:** machine-readable tables

## 1. Introduction

The Milky Way (MW) bulge exhibits a complex structure with composite stellar populations (Minniti 1996). Recent studies have found a composite Galactic bulge with at least two populations (metal-rich  $[\text{Fe}/\text{H}] \sim +0.3$  and metal-poor  $[\text{Fe}/\text{H}] \sim -0.4$ ) having different kinematics and spatial distributions (e.g., Rojas-Arriagada et al. 2017; Zoccali et al. 2017, and references therein). The different structures coexisting in the inner MW may be due to different formation mechanisms. But when we want to select the first born “primordial” populations, it is logical to concentrate on the most metal-poor stars. The RR Lyrae (RRL) pulsating variable stars are well-known tracers of old and metal-poor populations, often present in globular clusters, in the Galactic halo, and bulge (e.g., Catelan et al. 2004; Smith 2004). They are excellent distance and reddening indicators (Walker 1989; Longmore et al. 1990; Carney et al. 1995; Alcock et al. 1998; Bono et al. 2001; Catelan et al. 2004; Sollima et al. 2006), and

are found in large numbers in the Galactic bulge (Soszyński et al. 2011; Dékány et al. 2013; Pietrukowicz et al. 2015). Even though the dominant population of the bulge is metal-rich (e.g., Zoccali et al. 2017, and references therein), its RRL stars are metal-poor, with mean  $[\text{Fe}/\text{H}] \approx -1$  dex (Smith 1984; Butler et al. 1976; Rodgers 1977; Walker & Terndrup 1991). They also seem to have primordial He abundance, similar to old Galactic globular clusters (Marconi & Minniti 2018). As expected, the bulge RRL stars also have hot kinematics, with low rotation and large velocity dispersion (e.g., Gratton 1987; Minniti 1996; Kunder et al. 2016). These RRL are therefore ideal probes to study the oldest stars in the inner regions of our Galaxy, close to the Galactic Center, where large differential extinction and crowding complicate the discrimination of different populations. Indeed, previous near-infrared (NIR) searches managed to find only a dozen RRab in the Galactic Center region (Minniti et al. 2016; Dong et al. 2017).

The VISTA Variables in Via Láctea (VVV) is a deep NIR survey carried out to identify variable stars in some of the most crowded and reddened regions of the Galaxy (Minniti et al. 2010; Saito et al. 2012). The survey includes the region of the

\* Based on observations taken with ESO telescopes at Paranal Observatory under programme IDs 179.B-2002.

Galactic Center, enabling the discovery of RRL variables in this region (Minniti et al. 2016), which are the focus of this study, where we present a new catalog of about a thousand RRL type ab (RRab) stars. This is a vast improvement in the number of known RRL upon previous studies (Minniti et al. 2016; Dong et al. 2017), and allows the study and characterization of the RRab population in the most central region of our Galaxy.

This paper is organized as follows. Section 2 describes the selection of the RRL stars, while Section 3 presents color-magnitude diagrams (CMDs) for our sample. Section 4 discusses the reddening and extinction estimates, and in Section 5 we discuss the period distribution and inferred metallicity estimates. The computed distances are provided in Section 6, which likewise includes a discussion concerning the sample’s spatial distribution, and in Section 7 we discuss the total density of the old and metal-poor population. Section 8 presents the proper motions (PMs) and the derived kinematics for the sample. Section 9 presents a comparison with other RRL populations. Finally, the conclusions are summarized in Section 10.

## 2. The VVV Survey RR Lyrae Selection

The VVV survey has been acquiring data since early 2010 using the VISTA InfraRed Camera (VIRCAM) mounted on the VISTA telescope in Cerro Paranal Observatory, Chile. VIRCAM is a 16-detector array of  $2048 \times 2048$  pixels each, with a pixel scale of 0.34 arcsec/pix (Dalton et al. 2006; Emerson & Sutherland 2010). The VVV observation schedule includes single-epoch photometry in *ZYJH* bands and a variability campaign in the  $K_s$  filter (Minniti et al. 2010). At each sky pointing, six dithered images called pawprints are acquired and then combined in a single “tile,” a mosaic where the large gaps between the pawprint detectors are filled. The entire VVV observations comprise 348 tiles, 194 tiles in the bulge and 152 in the disk area (Saito et al. 2012), each covering a field of view (FoV) of  $\sim 1.6$  sq degrees. Public aperture photometric catalogs for all the regions covered by the VVV survey are provided by the Cambridge Astronomical Survey Unit (CASU) using the VISTA Data Flow System Pipeline (Emerson et al. 2004; Irwin et al. 2004); however, in this work we focus on the innermost region of the Galaxy, where the crowding is so severe that PSF modeling is mandatory. Accordingly, the photometric reduction of each detector was carried out using the DAOPHOT II/ALLSTAR package (Stetson 1987) and the CASU catalogs were used to calibrate our photometry into the VISTA system by means of a simple magnitude shift using several thousand stars in common (Contreras Ramos et al. 2017). We applied this procedure separately on each detector of the VVV tiles surrounding the inner 100 arcmin across the Galactic Center, comprising tiles b304, b305, b318, b319, b320, b332, b333, b334, b346, b347, and b348. The reduced data include  $\sim 100$  epochs spanning six years (2010–2015) of observations.

Searching for relatively faint and low-amplitude variable stars in the center of the MW, as is the case of RRL stars, is hampered by extremely high extinction and crowding. However, the dust in this central region of our Galaxy is far from being homogeneously distributed (see, for example, Gonzalez et al. 2012; Alonso-García et al. 2017, and references therein), meaning that RRL stars residing in the very inner bulge show quite different apparent magnitudes (but very similar reddening-corrected ones).

Thus, those RRL stars less affected by extinction are detectable using VVV data, as revealed in our initial search (Minniti et al. 2016).


RRL stars are commonly found as two main types: RRab, which pulsate in the fundamental mode and have longer periods and typically higher amplitudes than the second type, RR type c (RRc), which are first overtone pulsators. Part of the difficulty of finding RRL stars in the NIR is that their amplitudes of variation are smaller than those seen in the optical, typically  $A(K_s) \sim 1/3A(V)$  (e.g., Angeloni et al. 2014; Navarrete et al. 2015, 2017; Braga et al. 2018). The identification of RRc stars is even trickier because their light curves are rather sinusoidal, which is not very different from the morphology found in short-period eclipsing binaries. For these reasons, we focus our search on RRab only, which feature a unique sawtooth light curve, longer periods, and higher amplitudes. Their identification was made using visual inspection and two relatively new automated RRab classifiers, one described in Elorrieta et al. (2016) and a new algorithm especially developed for this work (see the Appendix). Their periodicities were obtained by means of the analysis of the variance statistic (Schwarzenberg-Czerny 1989) in the RRL period range. Representative light curves of our RRab sample can be seen in Figure 2 of Minniti et al. (2016).

Figure 1 shows the map of the RRab candidates found within 100 arcmin from the Galactic Center. As can be seen, most of the detected RRL stars are projected in regions of the inner bulge where the extinction is not extremely high. In fact, from the figure it is clear the dearth of RRL in the region  $|b| \lesssim 1^\circ$  is due to the heavy extinction close to the Galactic plane. The final catalog presented here includes 960 objects in total, some of which have been previously discovered (Soszynski et al. 2014; Minniti et al. 2016). For each RRL, we provide in Table 1 galactic coordinates, NIR magnitudes in the five VVV filters, and relative PMs with their corresponding statistical errors. In Table 2 we include extinction parameters, distance values, periods,  $K_s$  amplitudes, and metallicities.

The top panel of Figure 2 shows the  $K_s$ -band magnitude versus Galactic latitude for the 960 RRL candidates, illustrating the effect of extinction. As one approaches the Galactic plane, at latitudes  $|b| \lesssim 1^\circ$  the mean  $K_s$ -band magnitudes get fainter, until they fall beyond our detection limit at  $K_s \sim 16$  mag. Again, the lack of RRab in the Galactic plane is caused by heavy extinction, and our sample is severely incomplete in the regions where  $E(J - K_s) \gtrsim 3$  mag. The middle panel of Figure 2 shows the same figure using the reddening-corrected Wesenheit magnitudes, defined as  $W_{K_s} = K_s - 0.428 \times (J - K_s)$ , and discussed in detail in Section 4. When the extinction is taken into account, these Wesenheit magnitudes depend on the RRab periods, as shown in the bottom panel of Figure 2. After applying a  $3\sigma$  clipping to the  $P - W_{K_s}$  distribution, this diagram allows us to make a cut in order to separate Galactic Center RRL stars from the foreground and background counterparts. The diagram singles out the presence of 5 distant RRab and 25 RRab located in front of the Galactic bulge. We also note that there is one star that has no  $J$ -band photometry.

## 3. The CMD

The observed  $K_s$  versus  $J - K_s$  CMD of the VVV tiles b333 and b319, containing the Galactic Center and part of the area studied here, is shown in Figure 3. This region contains  $\sim 7$  million point sources that are plotted as a density diagram. Our sample of 959



**Figure 1.** Location in Galactic coordinates of the RRL candidates stars within a  $4 \times 4.5$  sq deg map surrounding the Galactic Center. The red dots mark the individual positions of the 960 RRL candidates located within 100 arcmin of the Galactic Center discovered with the NIR observations of the VVV Survey. At the distance of the Galactic Center, the scale is  $1^\circ \simeq 150$  pc. North galactic pole is up, and positive galactic longitudes are to the left.

**Table 1**  
RR Lyrae Sample Photometric Observations and Relative Proper Motions


ID	$L$ (deg)	$B$ (deg)	$K_s$	$H$	$J$	$Y$	$Z$	$\mu_l \cos(b)$ (mas yr $^{-1}$ )	$\epsilon \mu_l \cos(b)$	$\mu_b$ (mas yr $^{-1}$ )	$\epsilon \mu_b$
b333_312_65743	359.6790	-0.1952	14.284	15.886	18.170	20.360	99.999	6.753	2.65	-1.074	1.82
b333_616_55278	0.0533	-0.3683	15.626	16.817	19.066	20.816	99.999	-0.185	1.40	10.031	1.95
b333_614_37068	0.1636	0.2193	15.269	16.452	18.343	20.005	20.990	-2.541	1.49	0.974	1.89
b334_203_80042	0.2916	-0.0445	15.446	16.660	18.627	20.225	99.999	-2.819	1.74	-2.863	1.87
b333_314_84844	359.9308	0.3105	15.893	17.241	19.686	99.999	99.999	2.075	2.75	6.733	3.94
b333_414_58214	0.1097	0.3636	15.804	16.880	18.868	99.999	20.894	1.461	1.78	-1.711	1.94
b333_414_55144	0.0988	0.3692	14.639	15.866	18.003	99.999	99.999	5.190	1.63	-2.867	1.88
b334_104_33007	0.3240	-0.3333	15.287	16.634	18.737	20.503	99.999	4.708	2.24	3.267	2.41
b333_414_42332	0.1334	0.3928	15.344	16.360	17.934	19.433	21.403	-4.462	1.64	4.458	1.78
b333_414_8444	0.2182	0.4549	15.531	16.671	18.597	20.211	99.999	0.093	3.03	-5.529	3.03
b334_302_33289	0.3343	0.4071	15.581	16.666	18.613	99.999	99.999	6.577	2.61	-6.689	2.26
b305_209_26117	359.5813	-1.5817	14.636	14.895	15.527	16.342	16.983	3.713	1.04	-2.362	1.16

(This table is available in its entirety in machine-readable form.)

RRL candidates (one RRAb has no  $J$ -band photometry) is overlaid on the CMD as black dots. For comparison, the sample of  $\sim 1000$  outer bulge RRAb from Gran et al. (2016) is also plotted (white dots). The latter is almost unreddened because the outer bulge fields have  $E(J - K_s) < 0.2$  mag, and the former is a more differential reddened population that lies along the direction of the reddening vector, as expected. The reddening vector shown in Figure 3 with the black arrow has a slope  $\Delta K_s / \Delta(J - K_s) = 0.428$ , following the new results of Alonso-García et al. (2017) obtained using VVV data in the inner region of the Galaxy.

As they are all at approximately the same distance, we have used the location of the RRAb stars in the CMD to compute the total-to-selective extinction ratio at the Galactic Center, which in the presence of differential reddening should follow the slope  $\Delta K_s / \Delta(J - K_s)$ . The left panel of Figure 4 shows the CMD in the Galactic Center region, where we use a Hess density diagram for the RRAb sample and blue dots to separate the foreground and background objects from the rest of the population. From a simple least-squares fitting, we obtained  $A_{K_s} / E(J - K_s) = 0.438 \pm 0.016$  mag, in very good agreement with the values





**Figure 2.** Top:  $K_s$ -band magnitude vs. Galactic latitude for the 959 RRL candidates located within 100 arcmin of the Galactic Center. The effect of higher extinction closer to the Galactic plane is clear. Center: extinction-corrected (Wesenheit)  $W_{K_s}$ -band magnitude vs. Galactic latitude of the sampled variables. Bottom:  $W_{K_s}$ -band magnitude vs. period. This diagram allows us to make a cut in order to separate Galactic Center RRL (red dots) from foreground/background RRL and other contaminating objects (30 blue dots).

**Table 2**  
Measured Stellar Parameters


ID	$E(J - K_s)$	$A_{K_s}$	$(m - M)_0$	$D_{\odot}$ (kpc)	$P$ (days)	$A(K_s)$	[Fe/H] (dex)
b333_114_40405	4.101	1.755	14.792	9.086	0.78060	0.359	-2.21
b333_616_55278	3.270	1.400	14.751	8.915	0.62488	0.324	-1.66
b333_614_37068	2.904	1.243	14.539	8.087	0.61856	0.287	-1.64
b334_203_80042	3.011	1.289	14.213	6.960	0.40815	0.307	-0.62
b333_314_84844	3.623	1.551	14.726	8.814	0.54964	0.383	-1.35
b333_414_58214	2.894	1.238	14.609	8.351	0.40315	0.286	-0.59
b334_104_33007	3.280	1.404	14.401	7.590	0.62128	0.294	-1.65
b333_414_42332	2.420	1.036	14.630	8.435	0.52009	0.270	-1.21
b333_414_8444	2.896	1.240	14.542	8.098	0.48704	0.395	-1.05
b334_302_33289	2.862	1.225	14.591	8.284	0.48048	0.358	-1.02
b305_209_26117	0.721	0.309	14.698	8.703	0.54339	0.298	-1.32

(This table is available in its entirety in machine-readable form.)

obtained by Alonso-García et al. (2015, 2017). The effect of the extinction is taken into account in the right panel of Figure 4, which shows the  $W_{K_s}$  versus  $J - K_s$  CMD. Note that the mean

number of observations (different frames where  $K_s$  and PMs are measured) is  $N = 82$  in the  $K_s$ -band, with a range from  $N = 52$  to  $N = 104$ . Therefore, the mean magnitude errors are negligible for





**Figure 3.** VVV  $(J - K_s)$  vs.  $K_s$ -band color-magnitude diagram for the Galactic Center region (tiles b333 and b319) shown as a Hess density diagram (Minniti et al. 2016). The sample of  $\sim 1000$  outer bulge RRL from Gran et al. (2016) is plotted with white dots for comparison. Our 959 RRL candidates located within 100 arcmin of the Galactic Center are plotted as black dots. The reddening vector with slope  $\Delta K_s / \Delta (J - K_s) = 0.428$  is shown with the arrow (Alonso-García et al. 2017).

the  $K_s$ -band, because we are using the means of these multiple epochs, resulting in  $\sigma K_s = 0.0039 \pm 0.0009$  mag. However, fewer epochs (typically 2–4) are available for the other passbands. Therefore, the errors in the  $J$ -band, for example, are almost an order of magnitude higher in the mean,  $\sigma J = 0.020 \pm 0.019$  mag for the sample stars. Even worse, extinction affects the shorter passbands severely, and the mean errors are much larger in the mean  $\sigma Y = 0.04$  mag for the whole sample. These errors, however, have no major effect on the appearance of the CMDs shown in this work, which are rather dominated by extinction effects.

#### 4. Reddenings and Extinctions

The NIR reddening law in the Galactic Center region has been the subject of a variety of studies (see Matsunaga et al. 2009; Alonso-García et al. 2015; Majaess et al. 2016; Nataf et al. 2016; Alonso-García et al. 2017, and references therein). What matters for the present study is the slope of this reddening law in the NIR, which we take from the work recently published by Alonso-García et al. (2017),  $A_{K_s} / E(J - K_s) = 0.428$ . This allows us to define the reddening-corrected Wesenheit individual magnitudes,  $W_{K_s} = K_s - 0.428 \times (J - K_s)$  mag, for the RRL sample. The final distribution of the  $W_{K_s}$  mag magnitudes, shown in Figure 5, exhibits a strong peak at the expected location of the Galactic Center.


Moreover, RRL stars are excellent reddening indicators because in the NIR their temperatures are restricted to a very narrow range in the instability strip. Therefore, it is possible to use their observed colors to estimate reddenings along their lines of sight. Based on the data from Navarrete et al. (2015), we found that the intrinsic color of RRL can be assumed to be  $\Delta (J - K_s) = 0.17 \pm 0.03$  mag. Thus, the individual reddening values listed in Table 1 were obtained by the relation

$E(J - K_s) = (J - K_s) - 0.17$  mag. We inspect the spatial distribution of the bluest and reddest RRL in the sky region surrounding the Galactic Center. Figure 6 shows this map for the RRL candidates, color-coded by their respective reddenings. The darkest points represent the most reddened objects with  $E(J - K_s) > 3.5$  mag (corresponding to  $A_{K_s} \gtrsim 1.5$  mag), outlining the most heavily reddened regions close to the Galactic plane. On the other extreme, the lightest yellow dots show the location of the bluest objects with  $E(J - K_s) < 1.0$  mag, marking the less reddened regions (corresponding to  $A_{K_s} < 0.43$  mag). The bluest/reddest RRL stars of our sample have colors  $J - K_s \sim 0.61/4.3$  mag, equivalent to  $A_{K_s} \sim 0.19/1.75$  mag. In order to further characterize our sample, we have also examined the magnitude dependence on the RRL periods. Figure 7 (left panel) shows the  $Z$ -band magnitude versus period in days for our RRL candidates located within 100 arcmin of the Galactic Center. This figure clearly illustrates the damaging effects of reddening in the passbands at shorter-wavelengths, while this dependence is alleviated for the  $K_s$ -band magnitude (right panel). Even though the scatter due to reddening is still considerable, there is a hint of a period-luminosity (PL) relation seen in the  $K_s$ -band diagram. This relation becomes much tighter using the reddening-corrected Wesenheit magnitudes  $W_{K_s}$ , as shown in the bottom panel of Figure 2


#### 5. The Period Distribution and Metallicities

Figure 8 shows the  $K_s$  amplitude versus period in days (a.k.a. Bailey diagram) for the candidate RRL located within 100 arcmin from the Galactic Center, compared with (1) the  $I$ -band Bailey diagram sampling  $\sim 27,000$  OGLE RRL stars located in the middle bulge from Pietrukowicz et al. (2015), (2) the outer bulge sample of  $\sim 1000$  RRL stars from Gran et al. (2016) in the  $K_s$  band, and (3) the outer halo sample of  $\sim 10,000$  RRL stars published by Torrealba et al. (2015) in the  $V$  band. As can be seen in the figure, the three bulge distributions are similar, and no major differences are found. The ridge lines of the Oosterhoff I and II (OoI, OoII) population from Navarrete et al. (2015), Kunder et al. (2013), and Zorotovic et al. (2010) for the  $K_s$  VI bands, respectively, are indicated, showing that the bulk of Galactic bulge RRL stars, independent of their location, do not follow the same locus of the MW globular clusters, and in general are shifted to shorter periods for a given amplitude. It is likely that this period shift is because bulge RRL stars are in general more metal-rich than their counterparts in globular clusters. Similar results were obtained by Kunder & Chaboyer (2009) from their study of bulge RRL stars using MACHO data. On the other hand, outer halo RRL stars nicely match the OoI line, like the majority of relatively metal-rich globular cluster RRL stars. Quantitatively, the mean periods of the different samples shown in Figure 8 are:  $P_{ab} = 0.5446 \pm 0.0025$  days for our sample of Galactic Center RRL,  $P_{ab} = 0.5571 \pm 0.0005$  days for the OGLE bulge sample from Pietrukowicz et al. (2015),  $P_{ab} = 0.5634 \pm 0.0028$  days for the outer bulge RRL from Gran et al. (2016), and  $P_{ab} = 0.5751 \pm 0.0007$  days for the halo RRL sample from Torrealba et al. (2015), where the error is the standard error of the mean. These mean periods clearly decrease with decreasing distance to the Galactic Center, probably indicating that the innermost RRL stars are more metal-rich in the mean.


Our Bailey diagram can also be compared with the OGLE Bailey diagram to estimate the completeness of our RRL



**Figure 4.** Left: VVV  $K_s$ -band vs.  $(J - K_s)$  color-magnitude diagram for the 959 RRL candidates located within 100 arcmin of the Galactic Center. RRL located at the Galactic Center distance are plotted as a Hess diagram, while blue dots are foreground/background objects. Right: same as the left panel but using the extinction-corrected (Wesenheit)  $W_{K_s}$ -band magnitude. This was computed assuming a slope of  $\Delta K_s/\Delta (J - K_s) = 0.438$  from a fit shown by the red arrow in the right panel. This slope agrees within the errors with the slope measured by Alonso-García et al. (2015, 2017).



**Figure 5.** Extinction-corrected (Wesenheit)  $W_{K_s}$ -band magnitude distribution for the 959 RRL candidates located within 100 arcmin of the Galactic Center, after eliminating foreground/background stars. According to the magnitude distribution, the vast majority of the RRL candidates are located at the Galactic Center region.




**Figure 6.** Map of the candidate RRL, distinguished by color according to their respective  $E(J - K_s)$ . The darkest dots are RRL with colors  $J - K_s > 4$  ( $A_{K_s} \sim 1.7$  mag), outlining the most reddened regions. On the other hand, the lightest yellow dots show RRL with colors  $J - K_s < 1.0$ , indicating the less reddened regions. The black circle shows the area analyzed by Minniti et al. (2016).


sample. The NIR light curve amplitudes are lower than those in the optical by more than a factor of 2. In particular, we have very few ( $<1\%$  of the total sample) low-amplitude RRab in our sample with  $A(K_s) < 0.15$  mag, while the OGLE sample still includes about 10% of their low-amplitude population with  $A(J) < 0.3$  mag. From this simple comparison we can conclude that our sample is 10% incomplete, because we are missing low-amplitude RRab variable stars. The period distribution for our RRL sample is shown in the left panel of Figure 9.

The RRab periods can be used to estimate individual metallicities for RRab stars, taking, for example, the RRab stars of Layden (1994) as calibrators. Different authors have used these relations, which we have also applied for the VVV survey RRab in


the plane (Minniti et al. 2017b). Here we use the relations of Feast et al. (2010):  $[\text{Fe}/\text{H}] = -5.62 \log P_{ab} - 2.81$ , and Yang et al. (2010):  $[\text{Fe}/\text{H}] = -7.82 \log P_{ab} - 3.43$ . Using these relations, we obtain a mean metallicity of  $[\text{Fe}/\text{H}] = -1.30 \pm 0.01$  ( $\sigma = 0.33$ ) dex, and  $-1.33 \pm 0.02$  ( $\sigma = 0.46$ ) dex, respectively. The right panel of Figure 9 shows the metallicity distribution obtained using the relation of Feast et al. (2010), which is adopted for consistency with our previous work (Minniti et al. 2016, 2017b), and because it gives a smaller dispersion. The individual metallicities estimated using the periods are uncertain (Feast et al. 2010; Yang et al. 2010), and spectroscopic measurements are highly desirable. While the mean is metal-poor, the derived



**Figure 7.** Left: Z-band magnitude vs. period in days for the RRL candidates located within 100 arcmin of the Galactic Center (red points). Right:  $K_s$ -band magnitude vs. period in days for our sample. The blue points represent foreground/background objects.



**Figure 8.** Amplitude vs. period in days (Bailey diagram) for bulge RRL from different samples. From left to right we show the Bailey diagram for our candidate RRL located within 100 arcmin from the Galactic Center; all the OGLE RRab stars from Pietrukowicz et al. (2015); the outer bulge sample of 1000 RRab stars from Gran et al. (2016); and the halo RRab from Torrealba et al. (2015). The ridge lines of the Oosterhoff I and II populations are indicated in the panels (Zorotovic et al. 2010; Kunder et al. 2013; Navarrete et al. 2015) for comparison. Note that the OGLE and halo samples (2nd and 4th panels) show the amplitudes in the  $I$  band and  $V$  band respectively, while the present sample and outer bulge sample from the VVV survey show the amplitudes in the  $K_s$  band. This figure clearly shows that the Galactic Center RRL are more extreme than a typical OoI population.



**Figure 9.** Left: period distribution for our RRab sample compared with OGLE data (black Pietrukowicz et al. 2015) and outer halo RRab stars (green Torrealba et al. 2015). There appears to be a mild gradient in period but the differences are small. Right: same as the left panel but for the metallicity distribution. The  $[Fe/H]$  has been derived using the relation of Feast et al. (2010). The mean and the corresponding standard error of the period and metallicity for our RRab sample are indicated at the top of the panels.



metallicity distribution is very wide, and allows us to confirm that the sample contains both metal-poor and metal-rich RRab, consistent with the Bailey diagram that shows a mixture of Oosterhoff type I and II populations. In comparison, the periods for the OGLE sample yield an almost identical mean metallicity,  $[\text{Fe}/\text{H}] = -1.35$  dex using the relation of Feast et al. (2010). Also, the outer bulge population and outer halo RRab sample give similar metallicities ( $[\text{Fe}/\text{H}] = -1.38$  dex and  $[\text{Fe}/\text{H}] = -1.44$  dex). There appears to be a small gradient in the mean abundances, as RRL in the outer halo are more metal-poor than RRL in Galactic Center, but the systematics are big and the difference is within the errors. Moreover, we do not observe a significant metallicity gradient in the RRab population within  $1.5 > R_G > 0$  kpc.

## 6. Distances

In this section we derive distances for our RRL sample, following the procedures described by Gran et al. (2016), and Minniti et al. (2016). Briefly, we use the extinction law recently derived by Alonso-García et al. (2017), the PL calibration of Muraveva et al. (2015),  $M_{K_s} = -2.53 \log P_{ab} - 0.95 + 0.07 [\text{Fe}/\text{H}]$ , and we adopted  $[\text{Fe}/\text{H}] = -1.3$  dex as a good approximation of the individual metallicities (see Section 5). Even if the estimated abundance for our RRL stars is highly uncertain, the PL relation shows a very mild dependence on it. We also note that the impact in the computed distances due to the reduced number of *J*-band epochs is negligible. Figure 10 shows the distance distribution for the candidate RRL, separated by Galactic latitude, compared with the outer bulge RRab from Gran et al. (2016). This comparison is appropriate because all these samples have similar systematics, coming from the NIR VVV search for RRL. The median distance and the corresponding errors (standard error) of each subsample are indicated (from bottom to top: the outer bulge and the Galactic Center are negative and positive latitude, respectively). The figure shows how concentrated the Galactic Center RRL are: their dispersion is much smaller than the  $\sigma$  of the RRab located in the outer bulge. The median distance for the complete inner sample of 929 RRab stars is  $D = 8.052 \pm 0.024$  kpc ( $\sigma = 0.73$ ) and the median distance for the RRab sample of Gran et al. (2016), adopting  $A_{K_s}/E(J - K_s) = 0.689$  from Cardelli et al. (1989) and a mean metallicity  $[\text{Fe}/\text{H}] = -1.382$ , is  $D = 8.067 \pm 0.06$  kpc ( $\sigma = 1.85$ ). These distances agree within the errors of the distributions.

## 7. Densities

In this section we derive the number density for our RRL sample. This is important because it can give us an idea of how relevant the old and metal-poor population in the Galactic Center region is, where other populations (of young and intermediate age) are also present. In order to have a well-known example for comparison, we adopt the numbers corresponding to the giant globular cluster (or dwarf galaxy nucleus)  $\omega$  Centauri, for which Navarrete et al. (2015) presents a complete catalog of RRL stars using similar NIR data obtained with the VIRCAM at the VISTA telescope. Although the mean metallicities of  $\omega$  Centauri and bulge RRL stars are comparable, the overall color distribution of the HB stars in the central region of the Galaxy is not known. Therefore, we cannot guarantee that the bulge HB stars resemble  $\omega$  Centauri HB stars because of the difficulty of identifying HB stars in the


region. Thus, we stress that the comparison with  $\omega$  Centauri may be uncertain. According to Navarrete et al. (2015), there are  $N = 88$  RRab in this cluster (Navarrete et al. 2015), which has a mass of  $M = 4 \times 10^6 M_\odot$  (D’Souza & Rix 2013). That is, there are  $\sim 45,500 M_\odot$  per RRab in this globular cluster.

Figure 11 shows the RRab number counts as a function of projected Galactocentric distance in degrees (left panel), and across a narrow strip  $-0.3 < l < 0.3$  deg perpendicular to the Galactic plane (right panel). Based on the less reddened region sampled here we estimate a total density of  $\sim 1000$  RRL/sq deg at a Galactocentric distance of  $R_G = 1.6$  deg. Using the number ratio of RRab to total mass measured for  $\omega$  Centauri, this implies a total density of old and metal-poor stars of  $\sim 5 \times 10^7 M_\odot/\text{sq deg}$ . Considering the total region covered within 1.6 deg of the Galactic Center (the nuclear bulge of the MW), if we assume that the RRab volume density is constant, we obtain a total number of  $N \sim 13,000$  RRab inside this area. Obviously, this is an unreasonable assumption, but we take the flat density profile as a lower limit. Using this number we obtain a total mass of  $> 6 \times 10^8 M_\odot$  for the old metal-poor stars inside 1.6 deg. A more reasonable assumption in order to make an extrapolation into the inner regions is that the RRab keep their steep  $r^{-3.5}$  density profile, as observed by the MACHO and OGLE microlensing experiments in the Galactic bulge (Minniti et al. 1998; Pietrukowicz et al. 2015). Using this steep density profile we obtain a total number of  $N \sim 31,000$  RRab for the region with  $R_G < 1.6$  deg, from which we infer a total mass of  $> 1.4 \times 10^9 M_\odot$  for the old metal-poor stars. This is a large total mass that was obtained under two important assumptions: an  $\omega$  Cen-like population, and an inner  $r^{-3.5}$  density profile. These assumptions may be incorrect, and this number may be off by a factor of a few. However, we state that this total estimated mass would not be wrong by an order of magnitude in either direction. In fact, we can also argue that it is a lower limit because our sample is incomplete (at least 10% incomplete due to the lack of low-amplitude RRab with  $A(K_s) < 0.15$  mag, as discussed above), even in the less reddened region at 1.6 deg from the Galactic Center that we used to zero-point our density law.


Even though it is relatively small compared with the total mass of the Galactic bulge ( $M_{\text{bulge}} = 2 \times 10^{10} M_\odot$ ; Valenti et al. 2016), this mass of  $> 1.4 \times 10^9 M_\odot$  is a sizable contribution to the total mass budget of the Galactic nuclear bulge inside  $R_G = 1.6$  deg. This mass indicates that the old and metal-poor population may have played a dominant role in the inner dynamics of the MW, and also in the formation of the nuclear star cluster that contains the central black hole (BH; Mastrobuono-Battisti & Capuzzo-Dolcetta 2012). In fact, the central BH weights a small fraction of the mass present today in old and metal-poor stars:  $M_{\text{BH}} \simeq 4.2 \times 10^6 M_\odot$  (Ghez et al. 2008; Gillessen et al. 2009). One important implication would be that if the central BH formed very early, at the same time as the RRab studied here, and from similar material, its angular momentum may have been very low when it formed. If it has not spun up significantly by accreting disk material, a non-rotating supermassive BH would explain the absence of a strong jet, which is present in other supermassive BHs at the centers of distant galaxies.

## 8. Proper Motions


In this section we discuss the PM measurements for our RRab sample. These PMs have been measured following the procedures of Contreras Ramos et al. (2017), where a detailed description is given. The PMs computed in the present work are



**Figure 10.** Distance distribution for the candidate RRL, separated by Galactic latitude. The distances were computed considering those RRL that most likely belong to the Galactic bulge (stars between 5 and 11 kpc), and the median values of each subsample are indicated (from bottom to top: the outer bulge and the Galactic Center are negative and positive latitude, respectively). The median distances agree within the sigmas of the distributions. This figure shows how concentrated the RRL are at the Galactic Center.



**Figure 11.** Left: raw candidate RRL number density vs. distance from the Galactic Center in degrees. We find only about 270 RRL within  $1^\circ$  of the Galactic Center; most of the sample stars are located between 60 and 100 arcmin from the Galactic Center. Right: RRL number density across a strip with  $-0.3 < l < 0.3$  deg perpendicular to the Galactic plane. The less reddened regions away from the plane are more complete, reaching an observed total density of  $\sim 1000$  RRL/sq deg at  $b = 1.6$  deg. However, the number density of RRL drops drastically with decreasing Galactic latitude, where the sample becomes very incomplete due to extinction.




**Figure 12.** Bottom left panel: VPD in Galactic coordinates  $\mu_l \cos(b)$  vs.  $\mu_b$  in  $\text{mas yr}^{-1}$  for the candidate Galactic Center RRab stars. Top right panel: similar VPD for the stars classified as foreground objects. Top left: galactic longitude PM distribution  $\mu_l \cos(b)$  in  $\text{mas yr}^{-1}$  for the candidate Galactic Center RRL. Bottom right: same as the top left panel, for the Galactic latitude PM distribution  $\mu_b$  in  $\text{mas yr}^{-1}$ .

relative to the bulge population, obtained using the red giant branch as astrometric reference stars. The PMs are more complicated to obtain than our previous works because of the high field density; however, we achieve a mean precision that suffices for our purposes of checking the validity of the sample and for adding an extra criterion for statistically discriminating foreground from distant RRab stars.

Figure 12 shows the PM results in Galactic coordinates  $\mu_l \cos(b)$  versus  $\mu_b$  in  $\text{mas yr}^{-1}$  for the candidate Galactic Center RRab stars. The total number of RRab with relatively good PM measurements (total statistical error  $< 3 \text{ mas yr}^{-1}$ ) is 816, where we have also excluded 5 objects for which the PMs were impossible to measure because of crowding, and the 30 RRL deemed to be foreground/background objects (see the bottom panel of Figure 2). The lower left panel presents the vector point diagram (VPD) for the RRab sample showing a symmetric distribution, indicating a spherically symmetric spatial distribution. In particular, the Galactic longitude PM distribution does not have larger width than the Galactic latitude PM distribution, as one would expect from a flattened distribution or from a rotating population. This effect is even clearer in the lower right and top left panels of Figure 12, where the PM distribution in the Galactic longitude and latitude  $\mu_l \cos(b)$  and  $\mu_b$  in  $\text{mas yr}^{-1}$ , are shown. No significant asymmetries are observed in either of these distributions. We measure  $\sigma_{\mu_l \cos(b)} = 3.562 \pm 0.09 \text{ mas yr}^{-1}$ ,

and  $\sigma_{\mu_b} = 3.539 \pm 0.09$  for  $N = 816$  sample RRab stars with good PM measurements. The median statistical error in our PM measurements is  $1.3 \text{ mas yr}^{-1}$ , and considering  $0.37 \text{ mas yr}^{-1}$  due to systematic errors (Contreras Ramos et al. 2017), yields a total error budget of  $1.35 \text{ mas yr}^{-1}$  in each PM coordinate. After subtracting these errors in quadrature, we obtain  $\sigma_{\mu_l \cos(b)} = 3.29 \text{ mas yr}^{-1}$ , and  $\sigma_{\mu_b} = 3.27 \text{ mas yr}^{-1}$ , respectively. This allows us to conclude that the PM dispersions are real, and that the RRab population in the Galactic Center region intrinsically has a high velocity dispersion. In order to compute the tangential velocity dispersion we use  $V_T (\text{km s}^{-1}) = 4.74 \times \text{PM}(\text{arcsec yr}^{-1}) \times d (\text{pc})$ . At the distance of  $D_{\text{gc}} = 8.0 \text{ kpc}$ , these quantities correspond to tangential velocity dispersions of  $\sigma_{V_l} = 125.0 \pm 3.4 \text{ km s}^{-1}$  and  $\sigma_{V_b} = 124.1 \pm 3.4 \text{ km s}^{-1}$ , respectively. Adopting a larger distance to the Galactic Center,  $D_{\text{gc}} = 8.3 \text{ kpc}$  (Dékány et al. 2013), results in  $\sigma_{V_l} = 129.7 \pm 3.5 \text{ km s}^{-1}$  and  $\sigma_{V_b} = 128.7 \pm 3.5 \text{ km s}^{-1}$ , respectively. This very high tangential velocity dispersion in both components (along Galactic longitude and latitude) is not surprising, in light of the new results of Zoccali et al. (2017) and Valenti et al. (2016), who found a central peak in the radial velocity dispersion in the inner bulge. In their innermost field at  $1.0 \text{ deg}$  from the Galactic Center, they measure  $\sigma_{\text{RV}} \simeq 125$  and  $145 \text{ km s}^{-1}$  for their metal-poor and metal-rich population of red giants, respectively. Note that they defined metal-poor as all the red





**Figure 13.** Left:  $W_{K_s}$ -band magnitude for the candidate Galactic Center RRL as a function of Galactic longitude proper motion  $\mu_l \cos(b)$  in  $\text{mas yr}^{-1}$ . Right: same as the left panel, but for  $W_{K_s}$ -band magnitude vs. the Galactic latitude proper motion  $\mu_b$  in  $\text{mas yr}^{-1}$ . The slopes of the distributions suggest that there is no dependence of PM with distance from the Galactic Center.

giants with  $-0.8 < [\text{Fe}/\text{H}] < -0.1$  dex, and the metal-rich giants with  $[\text{Fe}/\text{H}] > 0.0$ . Considering that the RRab measured here are the most extreme tail of the metal-poor distribution, it is then not surprising that they exhibit a large velocity dispersion. This is a useful comparison with the red giants, because unfortunately the radial velocities would be difficult to measure for our sample RRab.

Figure 13 plots the reddening-corrected Wesenheit magnitude dependence of the measured  $\mu_l \cos(b)$  and  $\mu_b$  PMs in  $\text{mas yr}^{-1}$ . This figure illustrates that there is not a strong dependence of PM with magnitude for the Galactic Center RRab. The computed slope in the case of  $W_{K_s}$  versus  $\mu_l \cos(b)$  also shows that there is no evidence for rotation, given that the brighter (closer) RRab have similar motions to the fainter (more distant) RRab stars.

While the PMs are not accurate enough to explore in detail the kinematics around the Galactic Center, they clearly show that there is no significant rotation of the population, where one would expect the PM in the Galactic longitude direction to be larger than the PM in the latitude direction. In fact, the PM distribution shown in Figure 12 is round, and we find that  $\sigma_{\mu_l \cos(b)} \simeq \sigma_{\mu_b}$ , indicating a fairly isotropic velocity distribution for our RRL sample. Our results are in good agreement with the study of radial velocities of BRAVA RRL stars at lower latitudes (Kunder et al. 2016).

We note that these RRL are fundamentally different in properties than the red giants in the same region, which have been found to be divided into two main populations: one metal-rich component with  $[\text{Fe}/\text{H}] \sim +0.3$  dex that follows a barred distribution, and another more metal-poor component with a mean  $[\text{Fe}/\text{H}] \sim -0.4$  dex that has a more spherical distribution, both populations showing not negligible net rotation (Zoccali et al. 2017). The RRL here are more metal-poor in the mean and show zero rotation, suggesting they are more consistent with the extension of the old halo populations into the inner regions (Minniti 1996). This is our second major

result: the RRL population in the Galactic Center region shows very high tangential velocity dispersion of  $\sigma_{V_l} = 125.0 \text{ km s}^{-1}$  and  $\sigma_{V_b} = 124.1 \text{ km s}^{-1}$  and not bulk rotation. The RRab are a very special population because they represent the oldest and most metal-poor tracers that one can measure. The additional fact that the net angular momentum of the old and metal-poor stellar population is zero is also important. This has profound implications for the formation of the central regions of the MW, and the Galactic Center in particular. This supports the dissipational collapse formation scenario of this region.

## 9. Discussion

Are the RRL in the vicinity of the Galactic Center noteworthy in any way? In this section we briefly discuss the comparison with the RRL populations of other regions and components of the MW. We address the similarities and differences with the RRab found in the Galactic disk, halo, and inner and outer bulge, in order to discern if the Galactic Center RRab come from a different population.

- (1) The inner bulge RRab. The microlensing surveys have discovered thousands of RRL stars in the inner bulge. In particular, the MACHO and OGLE surveys show similar distance and metallicity properties as our sample RRab (Alcock et al. 1998; Kunder & Chaboyer 2009; Pietrukowicz et al. 2015). These samples, however, do not overlap, because our RRL are located in very heavily reddened regions that were not accessible to the optical surveys.
- (2) The outer bulge RRab. Gran et al. (2016) found a thousand RRab in the bulge-halo transition region, with projected distances between 1.2 and 1.4 kpc from the Galactic Center. This sample also shows similar characteristics as our sample of Galactic Center RRab. For example, their Bailey diagram looks qualitatively very similar. The Galactic Center RRL, however, have slightly

shorter mean periods, suggesting a more metal-rich population than the outer bulge RRL.

- (3) The globular cluster  $\omega$  Cen. Navarrete et al. (2015) and Braga et al. (2018) recently completed the NIR census and analysis of the RRL population associated with  $\omega$  Cen, the most massive Galactic globular cluster.  $\omega$  Cen is the prototypical example of an Oosterhoff type I population. As a whole, the R Rab stars of our sample do not follow the same loci in the Bailey diagram as the R Rab that belong to this massive cluster. This is relevant because globular clusters may have been part of the building blocks of the inner MW (Capuzzo-Dolcetta 1993), but these building blocks should be different from  $\omega$  Cen-like globular clusters. However, Minniti et al. (2017a) have used VVV data in order to search for concentrations of R Rab that would belong to unidentified bulge globular clusters. They were able to identify a dozen candidate globular clusters, which of course need to be confirmed with radial velocity measurements, for example.
- (4) The VVV disk sample. Recently, Minniti et al. (2017b) presented a large sample of R Rab located in the plane of the MW. Collectively, these stars also have similar properties (like mean periods, amplitudes, and metallicities) to our R Rab in the Galactic Center.
- (5) The halo population. There are various recent surveys of the R Rab population in the Galactic halo (e.g., Torrealba et al. 2015, and references therein). The main differences that we find with the R Rab in the Galactic Center are that the outer halo R Rab have a higher relative number of OoII R Rab than our sample, and that the mean periods of the Galactic Center R Rab are shorter than the halo samples, indicating a more metal-rich population. The inner halo, however, seems to have global properties more similar to the Galactic bulge R Rab (Abbas et al. 2014).

## 10. Conclusions

The first RRL stars in the Galactic Center region were discovered and initially characterized by Minniti et al. (2016) and Dong et al. (2017). In this work we present an extended sample based on the VVV survey NIR photometry. We have discovered 960 R Rab candidate variable stars within 100 arcmin of the Galactic Center. We measured accurate positions, NIR mean magnitudes, colors, periods, amplitudes, and relative PMs, as well as estimate distances and rough metallicities for the whole sample.

The CMDs reveal the dramatic effects of high and variable extinction in the Galactic Center region. Adopting an intrinsic mean colors for the R Rab stars  $J - K_s = 0.17 \pm 0.03$  and judging from the measured NIR colors of the present RRL sample ( $0.6 < J - K_s < 4.27$ ), the reddening in this field ranges from  $E(J - K_s) = 0.44$  to 4.1, and the extinction ranges from  $A_{K_s} = 0.19$  to 1.75. We conclude that this is the major limitation for the discovery of RRL in this region. The reddening slope using the R Rab sample is measured to be  $A_{K_s}/E(J - K_s) = 0.438 \pm 0.016$ , which is consistent with recent measurements using VVV data (0.428; Alonso-García et al. 2017).

We measure individual distances using the VVV NIR photometry. The distance distribution shows that the bulk of the sample is located at the distance of the bulge, as expected.

We use the  $P - W_{K_s}$  relation to separate 25 RRL located outside of the bulge, at a distance within 5 kpc from the Sun. We also identify five distant RRL, located behind the bulge, at  $D \approx 12$  kpc, indicating that there are regions, where the extinction is less severe. The mean distance for our R Rab sample is  $D = 8.05 \pm 0.024$  kpc.

The observed density of R Rab variables in the Galactic Center region is very high (1000 RRL/sq deg at  $R = 1.6$  deg). This agrees with the extrapolation of the RR Lyr density profile from the bulge microlensing experiments into the inner regions (Minniti et al. 1998; Pietrukowicz et al. 2015), and implies that the old metal-poor populations make a significant contribution to the total mass budget in this region.

The mean period is  $P = 0.5446 \pm 0.0025$  days, yielding a mean metallicity of  $[Fe/H] = -1.30 \pm 0.01$  ( $\sigma = 0.33$ ) dex for the R Rab sample in the Galactic Center region. This sample is compared with the VVV RRL samples from the MW disk (Minniti et al. 2017b), middle bulge (Pietrukowicz et al. 2015), and outer bulge (Gran et al. 2016). These comparisons show reasonable agreement, and we do not find major differences among these different samples. However, the period distribution suggests a higher metallicity than the Oosterhoff type I population in the outer halo studied by Torrealba et al. (2015). The Bailey diagram also shows a predominant population that is more extreme than Oosterhoff type I RRL stars.

We measure PMs, firmly establishing the kinematics of the old and metal-poor population in the Galactic Center region. We find a very large velocity dispersion  $\sigma \approx 130$  km s $^{-1}$  for the R Rab variable stars, and no indication of significant rotation. The PM kinematics show that this specific bulge population may have formed by dissipational collapse, and the kinematics are also consistent with the scenario that the population may be an extension of the inner halo (Minniti 1996). The results are also consistent with the observed chemical history, as the bulge red giants seem to exhibit a rapid formation, judging from their measured chemical enrichment (e.g., O-enhancement found by Zoccali et al. 2008, and references therein). The measurement of radial velocities for the present sample would be very profitable for investigating the orbital dynamics of this old and metal-poor population. These RVs would also be useful to confirm or disprove a number of globular cluster candidates recently identified in the Galactic Center region as concentrations of R Rab (Minniti et al. 2017a).

The RRL population of the Galactic Center deserves more observations and theoretical modeling to see in what proportion it could come from disrupted globular clusters, what fraction of the population is inherent to the bulge, and what fraction is just passing by from the halo population. Indeed, these are useful prime targets for future spectroscopy, in order to investigate their 3D kinematics and chemical compositions. In addition, the present work underscores the need for the *WFIRST* space mission, which will be able to complete the RRL census that cannot be done from the ground in the most obscured regions. Without *WFIRST* this part of the Galaxy will remain hidden.

We gratefully acknowledge the use of data from the ESO Public Survey program ID 179.B-2002 taken with the VISTA telescope, and data products from the Cambridge Astronomical Survey Unit (CASU). Support for the authors is provided by the BASAL Center for Astrophysics and Associated Technologies (CATA) through grant PFB-06, and the Ministry for the

Economy, Development, and Tourism, Programa Iniciativa Científica Milenio through grant IC120009, awarded to the Millennium Institute of Astrophysics (MAS). D.M. and M.Z. acknowledge support from FONDECYT Regular grants No. 1170121, and 1150345, respectively. P.H. acknowledges financial support from FONDECYT regular grant 1170305. F.G. acknowledge support from CONICYT-PCHA Doctorado Nacional 2017-21171485 and Proyecto Fondecyt Regular 1150345. J.A.-G. acknowledges support by FONDECYT Iniciación 11150916. D.M. is also grateful for the hospitality of the Vatican Observatory. This research made use of Astropy, a community-developed core Python package for astronomy; Scikit-learn, NumPy, and matplotlib, a Python library for publication-quality graphics; and Aladin Sky Atlas, developed at CDS, Strasbourg Observatory, France, and TOPCAT.

## Appendix

### Positive-unlabeled (PU) Random Forest Classifier

PU learning refers to a case in which a set representing a known class (P) and a usually larger set containing unlabeled data (U) are used to train a semi-supervised classifier to retrieve the objects in U that are most similar to P. In this case P and U correspond to the RR Lyrae found in Gran et al. (2016) and the light curves of the bulge, respectively. We draw from Mordelet & Vert (2014) and train an ensemble of decision trees using a PU version of the bagging algorithm (a python implementation of the algorithm is available at [http://www.github.com/phuijse/bagging\\_pu](http://www.github.com/phuijse/bagging_pu)). To obtain features, the light curves are (1) folded, (2) aligned in phase, (3) normalized, and (4) interpolated to a 50-point regular grid using Gaussian Process (GP) regression. For U the period is found using mutual information (Huijse et al. 2018). Fifty additional feature foldings with twice the reported period are also included. The 100 GP features plus the amplitude and period are the input to the classifier. This differs from Elorrieta et al. (2016) in that the classifier uses the distribution of U (semi-supervision) and the features that characterize the light curves.

## ORCID iDs

Rodrigo Contreras Ramos  <https://orcid.org/0000-0001-7948-9731>

Dante Minniti  <https://orcid.org/0000-0002-7064-099X>

Felipe Gran  <https://orcid.org/0000-0002-9252-6899>

Manuela Zoccali  <https://orcid.org/0000-0002-5829-2267>

Javier Alonso-García  <https://orcid.org/0000-0003-3496-3772>

Pablo Huijse  <https://orcid.org/0000-0003-3541-1697>

María Gabriela Navarro  <https://orcid.org/0000-0002-1860-2304>

Álvaro Rojas-Arriagada  <https://orcid.org/0000-0003-0208-8854>

Elena Valenti  <https://orcid.org/0000-0002-6092-7145>

## References

- Abbas, M. A., Grebel, E. K., Martin, N. F., et al. 2014, *MNRAS*, **441**, 1230
- Alcock, C., Allsman, R. A., Alves, D. R., et al. 1998, *ApJ*, **492**, 190
- Alonso-García, J., Dékány, I., Catelan, M., et al. 2015, *AJ*, **149**, 99
- Alonso-García, J., Minniti, D., Catelan, M., et al. 2017, *ApJL*, **849**, L13
- Angeloni, R., Contreras Ramos, R., Catelan, M., et al. 2014, *A&A*, **567**, A100
- Bono, G., Caputo, F., Castellani, V., Marconi, M., & Storm, J. 2001, *MNRAS*, **326**, 1183
- Braga, V. F., Stetson, P. B., Bono, G., et al. 2018, arXiv:1802.03578
- Butler, D., Carbon, D., & Kraft, R. P. 1976, *ApJ*, **210**, 120
- Capuzzo-Dolcetta, R. 1993, *ApJ*, **415**, 616
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, **345**, 245
- Carney, B. W., Fulbright, J. P., Terndrup, D. M., Suntzeff, N. B., & Walker, A. R. 1995, *AJ*, **110**, 1674
- Catelan, M., Pritzl, B. J., & Smith, H. A. 2004, *ApJS*, **154**, 633
- Contreras Ramos, R., Zoccali, M., Rojas, F., et al. 2017, arXiv:1709.07919
- Dalton, G. B., Caldwell, M., Ward, A. K., et al. 2006, *Proc. SPIE*, **6269**, 62690X
- D'Souza, R., & Rix, H.-W. 2013, *MNRAS*, **429**, 1887
- Dékány, I., Minniti, D., Catelan, M., et al. 2013, *ApJL*, **776**, L19
- Dong, H., Schödel, R., Williams, B. F., et al. 2017, *MNRAS*, **471**, 3617
- Elorrieta, F., Eyheramendy, S., Jordán, A., et al. 2016, *A&A*, **595**, A82
- Emerson, J. P., Irwin, M. J., Lewis, J., et al. 2004, *Proc. SPIE*, **5493**, 401
- Emerson, J. P., & Sutherland, W. J. 2010, *Proc. SPIE*, **7733**, 773306
- Feast, M. W., Abedigamba, O. P., & Whitelock, P. A. 2010, *MNRAS*, **408**, L76
- Ghez, A. M., Salim, S., Weinberg, N. N., et al. 2008, *ApJ*, **689**, 1044
- Gillessen, S., Eisenhauer, F., Trippe, S., et al. 2009, *ApJ*, **692**, 1075
- Gonzalez, O. A., Rejkuba, M., Zoccali, M., et al. 2012, *A&A*, **543**, A13
- Gran, F., Minniti, D., Saito, R. K., et al. 2016, *A&A*, **591**, A145
- Gratton, R. G. 1987, *MNRAS*, **224**, 175
- Huijse, P., Estévez, P., Forster, F., et al. 2018, *ApJS*, **236**, 12
- Irwin, M. J., Lewis, J., Hodgkin, S., et al. 2004, *Proc. SPIE*, **5493**, 411
- Kunder, A., & Chaboyer, B. 2009, *AJ*, **138**, 1284
- Kunder, A., Rich, R. M., Koch, A., et al. 2016, *ApJL*, **821**, L25
- Kunder, A., Stetson, P. B., Catelan, M., Walker, A. R., & Amigo, P. 2013, *AJ*, **145**, 33
- Layden, A. C. 1994, *AJ*, **108**, 1016
- Longmore, A. J., Dixon, R., Skillen, I., Jameson, R. F., & Fernley, J. A. 1990, *MNRAS*, **247**, 684
- Majaess, D., Turner, D., Dékány, I., Minniti, D., & Gieren, W. 2016, *A&A*, **593**, A124
- Marconi, M., & Minniti, D. 2018, *ApJL*, **853**, L20
- Mastrobuono-Battisti, A., & Capuzzo-Dolcetta, R. 2012, in ASP Conf. Ser. 453, *Advances in Computational Astrophysics: Methods, Tools, and Outcome*, ed. R. Capuzzo-Dolcetta, M. Limongi, & A. Tornambè (San Francisco, CA: ASP), 237
- Matsunaga, N., Kawadu, T., Nishiyama, S., et al. 2009, *MNRAS*, **399**, 1709
- Minniti, D. 1996, *ApJ*, **459**, 175
- Minniti, D., Alcock, C., Alves, D., et al. 1998, in IAU Symp. 184, *The Central Regions of the Galaxy and Galaxies*, ed. Y. Sofue (Dordrecht: Kluwer), 123
- Minniti, D., Alonso-García, J., Braga, V., et al. 2017a, *RNAAS*, **1**, 16
- Minniti, D., Contreras Ramos, R., Zoccali, M., et al. 2016, *ApJL*, **830**, L14
- Minniti, D., Dékány, I., Majaess, D., et al. 2017b, *AJ*, **153**, 179
- Minniti, D., Lucas, P. W., Emerson, J. P., et al. 2010, *NewA*, **15**, 433
- Mordelet, F., & Vert, J.-P. 2014, *PaReL*, **37**, 201
- Muraveva, T., Palmer, M., Clementini, G., et al. 2015, *ApJ*, **807**, 127
- Nataf, D. M., Gonzalez, O. A., Casagrande, L., et al. 2016, *MNRAS*, **456**, 2692
- Navarrete, C., Catelan, M., Contreras Ramos, R., et al. 2017, *A&A*, **604**, A120
- Navarrete, C., Contreras Ramos, R., Catelan, M., et al. 2015, *A&A*, **577**, A99
- Pietrukowicz, P., Kozłowski, S., Skowron, J., et al. 2015, *ApJ*, **811**, 113
- Rodgers, A. W. 1977, *ApJ*, **212**, 117
- Rojas-Arriagada, A., Recio-Blanco, A., de Laverny, P., et al. 2017, *A&A*, **601**, A140
- Saito, R. K., Hempel, M., Minniti, D., et al. 2012, *A&A*, **537**, A107
- Schwarzenberg-Czerny, A. 1989, *MNRAS*, **241**, 153
- Smith, H. A. 1984, *PASP*, **96**, 505
- Smith, H. A. 2004, *RR Lyrae Stars* (Cambridge: Cambridge Univ. Press), 166
- Sollima, A., Cacciari, C., & Valentini, E. 2006, *MNRAS*, **372**, 1675
- Soszyński, I., Udalski, A., Szymański, M. K., et al. 2014, *AcA*, **64**, 177
- Soszyński, I., Dziembowski, W. A., Udalski, A., et al. 2011, *AcA*, **61**, 1
- Stetson, P. B. 1987, *PASP*, **99**, 191
- Torrebalba, G., Catelan, M., Drake, A. J., et al. 2015, *MNRAS*, **446**, 2251
- Valenti, E., Zoccali, M., Gonzalez, O. A., et al. 2016, *A&A*, **587**, L6
- Walker, A. R. 1989, *PASP*, **101**, 570
- Walker, A. R., & Terndrup, D. M. 1991, *ApJ*, **378**, 119
- Yang, S.-C., Sarajedini, A., Holtzman, J. A., & Garnett, D. R. 2010, *ApJ*, **724**, 799
- Zoccali, M., Hill, V., Lecureur, A., et al. 2008, *A&A*, **486**, 177
- Zoccali, M., Vazquez, S., Gonzalez, O. A., et al. 2017, *A&A*, **599**, A12
- Zorotovic, M., Catelan, M., Smith, H. A., et al. 2010, *AJ*, **139**, 357